

Document made available under the Patent Cooperation Treaty (PCT)

International application number: PCT/US2004/032341

International filing date: 01 October 2004 (01.10.2004)

Document type: Certified copy of priority document

Document details: Country/Office: US
Number: 60/507,514
Filing date: 02 October 2003 (02.10.2003)

Date of receipt at the International Bureau: 18 May 2008 (18.05.2008)

Remark: Priority document submitted or transmitted to the International Bureau in compliance with Rule 17.1(a) or (b)



World Intellectual Property Organization (WIPO) - Geneva, Switzerland
Organisation Mondiale de la Propriété Intellectuelle (OMPI) - Genève, Suisse



THE UNITED STATES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME:

UNITED STATES DEPARTMENT OF COMMERCE

United States Patent and Trademark Office

May 18, 2008

THIS IS TO CERTIFY THAT ANNEXED HERETO IS A TRUE COPY FROM THE RECORDS OF THE UNITED STATES PATENT AND TRADEMARK OFFICE OF THOSE PAPERS OF THE BELOW IDENTIFIED PATENT APPLICATION THAT MET THE REQUIREMENTS TO BE GRANTED A FILING DATE.

APPLICATION NUMBER: 60/507,514

FILING DATE: October 02, 2003

RELATED PCT APPLICATION NUMBER: PCT/US04/32341

THE COUNTRY CODE AND NUMBER OF YOUR PRIORITY APPLICATION, TO BE USED FOR FILING ABROAD UNDER THE PARIS CONVENTION, IS US60/507,514



Certified by

Under Secretary of Commerce
for Intellectual Property
and Director of the United States
Patent and Trademark Office

**U.S. PATENT AND TRADEMARK OFFICE
PROVISIONAL APPLICATION FOR PATENT COVER SHEET**

This is a request for filing a PROVISIONAL APPLICATION FOR PATENT
under 37 C.F.R. §1.53(b)(2)

Atty. Docket: CHANG184

| INVENTOR(S)/APPLICANT(S) | | | |
|---|------------------|----|--|
| LAST NAME | FIRST NAME | MI | RESIDENCE (CITY AND EITHER STATE OR FOREIGN COUNTRY) |
| CHANG | Young-Tae | | New York, New York |
| <input type="checkbox"/> Additional inventors are being named on separately numbered sheets attached hereto | | | |
| TITLE OF THE INVENTION (280 characters max) | | | |
| RECEPTOR SPECIFIC BINDER DISCOVERY USING INTRAMOLECULAR ACYL MIGRATION INDUCED DYNAMIC CARBOHYDRATE LIBRARY | | | |
| CORRESPONDENCE ADDRESS | | | |
| Direct all correspondence to the address associated with Customer Number 001444 , which is presently: BROWDY AND NEIMARK, P.L.L.C. 624 Ninth Street, N.W., Suite 300 Washington, D.C. 20001-5303 | | | |
| ENCLOSED APPLICATION PARTS (check all that apply) | | | |
| <input checked="" type="checkbox"/> Specification | Number of Pages | 22 | <input type="checkbox"/> Applicant claims small entity status. See 37 C.F.R. §1.27 |
| <input checked="" type="checkbox"/> Drawing(s) | Number of Sheets | 9 | <input type="checkbox"/> Other (specify) _____ |
| METHOD OF PAYMENT (check one) | | | |
| <input checked="" type="checkbox"/> Credit Card Payment Form PTO-2038 is enclosed to cover the Provisional filing fee of <input type="checkbox"/> \$160 large entity <input checked="" type="checkbox"/> \$80 small entity | | | |
| <input checked="" type="checkbox"/> The Commissioner is hereby authorized to charge filing fees and credit Deposit Account Number 02-4035 | | | |

The invention was made by an agency of the United States Government or under a contract with an agency of the United States Government.

☒ No ☐ Yes, the name of the U.S. Government agency and the Government contract number are:

Respectfully submitted,

BROWDY AND NEIMARK, P.L.L.C.

By: Anne M. Kornbau
Registration No.: 25,884

Date: October 2, 2003

AMK:ju

03916 U.S. PTO
60/507514

100203

RECEPTOR SPECIFIC BINDER DISCOVERY USING INTRAMOLECULAR ACYL MIGRATION INDUCED DYNAMIC CARBOHYDRATE LIBRARY

Field of the Invention

[0001] The present invention provides a highly efficient molecular evolution system using intramolecular acyl migration on a carbohydrate scaffold coupled with boronic acid as a selector.

Background of the Invention

[0002] Evolution through natural selection underlies much of current theory of the biological world, from the molecular to the species level. Much effort has been expended trying to realize evolutionary phenomena at the molecular level (Otto et al., 2002; Furlan et al., 2002; Karan et al., 2000; Rowan et al., 2002). Dynamic combinatorial chemistry enabled functional features of evolution such as a diversity generation system (mutations) and survival of the fittest (natural selection).

[0003] A dynamic combinatorial library can be visualized as a test of transient compounds that undergo a series of reversible equilibrium processes. Under thermodynamic control, the library members interconvert reversibly and continually. The action of the selector drives the system to

re-equilibrate, forming an amplified concentration of the molecule with the highest selector binding affinity.

[0004] Two key features of molecular evolution are the selection and amplification of a particular species in the dynamic combinatorial library (DCL). A DCL allows for a reversible equilibrium of library members in which a concentration change in one library member results in a new equilibrium distribution following Le Chatelier's principle.

[0005] To date, a number of DCLS have been prepared using such diverse chemical reactions as intermolecular transesterification, enzyme catalyzed peptide-bond exchange, imine bond exchanges of hydrazones or oximes, olefin metathesis, disulfide bond exchange, photoisomerization, hydrogen bond exchange, and metal-ligand coordination.

[0006] An ideal molecular evolution system requires an even distribution of the DCL components, an efficient selection method, and a non-destructive/continuous equilibrium in order to generate a "winning" binder for amplification.

[0007] As molecular evolution needs an effective molecular recognition event for amplification, it is now being evaluated as an efficient approach for identifying and preparing new host-guest systems (Ott et al., 2002; Furlan et al., 2002; Furlan et al., 2001).

[0008] Carbohydrate recognition plays an important role in many biological processes, particularly in cell-cell interactions and cell communications. A multitude of regulatory processes are mediated by carbohydrates as ligands for diverse proteins. However, the preparation of carbohydrate libraries by conventional methods has not achieved a satisfactory development because of numerous problems met with carbohydrate synthesis and automation protocols. Although dynamic combinatorial chemistry (DCC) can offer the complementary method to carbohydrate libraries, there have not been many attempts to generate dynamic combinatorial libraries (DCL) on a carbohydrate scaffold because of the complexity, and the selection method was not effective for successful amplification.

Summary of the Invention

[0009] It is an object of the present invention to overcome the aforesaid deficiencies in the prior art.

[0010] It is another object of the present invention to provide a molecular evolution system.

[0011] It is yet another object of the present invention to provide a molecular evolution system based on a carbohydrate scaffold

[0012] It is a further object of the present invention to produce a dynamic combinatorial library of dimers and oligomers having monomers connected by an ester bond using intramolecular acyl migration.

[0013] The present invention provides a highly efficient molecular evolution system using intramolecular acyl migration on a carbohydrate scaffold coupled with boronic acid as a selector.

[0014] In the examples described below, hexahydroxyl cyclohexane (inositol) was used for the intramolecular acyl migration system. In these molecules, each free hydroxyl group behaves as a nucleophile by attacking neighboring acyl groups, thus generating various regioisomers. Under basic conditions (pyridine/water), it was previously reported that benzoyl migration on *myo*-inositol (with 5 equatorial and one axial OH) generates an almost equi-molar amount of 9 regioisomers (Chung et al., 1995; Chung et al., 1996).

[0015] To maximize the geometric diversity between regioisomers, *chiro*-inositol was chosen as a DCL scaffold. *Chiro*-inositol has 2 vicinal axial-OH and 4 equatorial OH groups. Accordingly, 1, 4-dibenzoyl-*chiro*-inositol was synthesized (Khersonsky et al., 2002; Falshaw et al., 2000) and investigated for optimization of the DCL generation.

[0016] The 1,4-dibenzoyl-*chiro*-inositol was synthesized and subjected to a benzoyl migration. Under basic conditions, benzoyl migration on the inositol ring generated an almost equi-molar amount of 9 regioisomers. To minimize debenzoylation, anhydrous acetonitrile was used as the solvent with DBU as the base. The optimal condition consisted of 1 mg 1,4-dibenzoyl-*chiro*-inositol and 30 mL DBU [WHAT IS DBU?] in 1 mL acetonitrile. This yielded full migration within one hour at room temperature. Generation of the nine isomers was confirmed by mass spectrometry and UV analysis using LC-MS, as shown in Figure 1a and Figure 1b.

[0017] Boronic acids can be added to the mixture as a selector, since it is well known that boronic acids form five-membered cyclic esters preferably with 1,2-cis diols in sugar. Adding the boronic acid shifted the equilibrium in favor of one isomer over the others.

Brief Description of the Drawings

[0018] Figure 1 shows library members generated by acyl migration and the selected component upon treatment with phenyl boronic acid.

[0019] Figures 2a-2d show HPLC chromatograms of *chiro*-1(1,4)Bz.

[0020] Figure 3 illustrates synthesis of 3,4-di-O-benzoyl-*chiro*-inositol, compound 7.

[0021] Figure 4 shows the effect of phenyl boronic acid on the amplification of 3,4-di-O-benzoyl-*chiro*-inositol.

[0022] Figure 5 shows time-dependent amplification of 3,4-di-O-benzoyl-*chiro*-inositol.

[0023] Figure 6 shows synthesis of 3,6-di-O-benzoyl-*chiro*-inositol, compound 5.

[0024] Figure 7 shows LC-MS diagram of 3,4-di-O-benzoyl-*chiro*-inositol (7a) and mass data for 3,4-di-O-benzoyl-*chiro*-inositol (7b).

[0025] Figure 8 shows LC-MS diagram of 3,6-di-O-benzoyl-*chiro*-inositol (8a) and mass data for 3,6-di-O-benzoyl-*chiro*-inositol (8b).

[0026] Figure 9 shows a dynamic combinatorial library on an oligosaccharide scaffold.

[0027] Figure 10 shows intramolecular acyl migration of *chiro*-inositol dibenzoate.

Detailed Description of the Invention

[0028] Hexahydroxyl cyclohexane (inositol) was chosen to illustrate the intramolecular acyl migration model system of the present invention. In these molecules, each free hydroxyl group behaves as a nucleophile by attacking neighboring acyl

groups, thus generating various regioisomers. Under basic conditions (pyridine/water), it was previously reported that benzoyl migration on *myo*-inositol (with five equatorial and 1 axial OH) generates an almost equi-molar amount of nine regioisomers Chung et al., 1995; Chung et al., 1996).

[0029] The process of the present invention maximizes the geometric diversity between regioisomers by using *chiro*-inositol, which has 2 vicinal axial OH groups and 4 equatorial OH groups as a dynamic combinatorial scaffold. Accordingly, 1,4-dibenzoyl-*chiro*-inositol was synthesized (Khersonsky et al., 2002; Falshaw et al., 2000) and investigated for optimization of the dynamic combinatorial library generation. *Chiro*-inositol dibenzoate generates a total of nine regioisomers upon full equilibration, as shown in Figure 1. A series of bases, including pyridine, DMAP*, DABCO*, DIEA*, BEMP*, DBU* and NaOH in aprotic solvents (DMF, DMSO, CH₂CN) all with and without water were evaluated. **[PLEASE PROVIDE NAMES FOR THESE BASES]**. While several conditions induce benzoyl migration at a reasonable rate, it was found that increasing amounts of water accelerated the formation of the dynamic combinatorial library due to the increased polarity of the solvent, but also increased the extent of hydrolysis of the benzoyl group. In addition, the migration rate strongly depends on the concentration and pK_b of the base and

temperature. To minimize debenzoylation, anhydrous acetonitrile was used as the solvent, with DBU as a base. The optimal condition consisted of 1 mg of 1,4-dibenzoyl-*chiro*-inositol (compound 5) and 30 microL (79 eq) of DBU in 1 mL acetonitrile, which yielded full migration within one hour at room temperature. Generation of the nine isomers was confirmed by mass spectroscopy and UV analysis using LC-MS, as shown in Figure 2.

[0030] Figure 2 shows (a) HPLC chromatogram (C18 column: 4.6 x 150 mm, eluted with 18% acetonitrile in water of 3,4-di-*O*-benzoyl-*chiro*-inositol before base addition; (b) chromatogram of the DCL at one hour after addition of DBU with 8% of 3,4-di-*O*-benzoyl-*chiro*-inositol (c) chromatogram of the DCL at seven hours after addition of 2 eq of PhB(OH)₂ with 43% of 3,4-di-*O*-benzoyl-*chiro*-inositol; and (d) chromatogram of the DCL at seven hours after addition of PhB(OH)₂ with 82% of 3,4-di-*O*-benzoyl-*chiro*-inositol.

[0031] Since it is well known that boronic acids form five-membered cyclic esters preferably with 1,2-*cis*-diols in sugar systems (James et al., 1996; Wiecko et al., 1979; Eggert et al., 1999; Arimori et al., 2001; Arimori et al., 2002; Yang et al., 2001; Cabell et al., 1999; James et al., 1995), the present inventors envisioned that only one isomer out of the nine, namely, 3,4-dibenzoyl-*chiro*-inositol, which carries two

cis-vicinal diols, would have a higher binding affinity for two boronic acids, thus stopping further migration. Furthermore, it was also reported that the binding affinity of boronic acid with a sugar depends on the pH of the media. At pH lower than the pKa of boronic acid, *i.e.*, acidic conditions, coupling between the boronic acid and a sugar is disfavored, while the coupling is favored at pH over the pKa of boronic acid, *i.e.*, basic donations.

[0032] To test this hypothesis, 32 eq of phenyl boronic acid was added to the equilibrium mixture of the *chiro*-inositol dibenzoates in the presence of DBU. A dramatic equilibrium shift of mixtures toward 3,4-di-*O*-benzoyl-*chiro*-inositol, compound **7**, characterized as 3,4-di-*O*-benzoyl-*chiro*-inositol by a separate synthesis (Khersonsky et al., 2002; Falshaw, 2000), was demonstrated, as shown in Figure 2c. The isomer 3,4-di-*O*-benzoyl-*chiro*-inositol, initially 8% of the equilibrium mixture, as shown in Figure 2b, became the major constituent of the library (82%) in seven hours after treatment with DBU, demonstrating an enrichment factor of 10.3.

[0033] It was envisioned that the nine regioisomers generated from compound **5**, 3,6-di-*O*-benzoyl-*chiro*-inositol, would initially couple with one phenyl boronic acid at the moment one cis-diol in any component was exposed from benzoyl

migration. Following that, the mono-boronic acid-coupled component will undergo further acyl migration until a second cis-diol is exposed and trapped by yet another phenyl boronic acid, as shown in Figure 1. Benzoyl migration will terminate when two cis-diols of 3,4-di-O-benzoyl-*chiro*-inositol couple with two boronic acids, since free hydroxyl groups are no longer available. This effectively isolates this component from the equilibrating DCL, resulting in an accumulation of the "winning" binder. Thus, increased amounts of boronic acid will shift the equilibrium towards the selected compound, 3,4-di-O-benzoyl-*chiro*-inositol. With 1, 2, 6 and 32 equivalents of phenyl boronic acid, the final amount of 3,4-di-O-benzoyl-*chiro*-inositol was 25%, 43%, 67%, and 82%, respectively. However, amounts of phenyl boronic acid in excess of 32 equivalents did not further amplify 3,4-di-O-benzoyl-*chiro*-inositol. The final distribution of the regioisomers is "quenched" by treating the reaction mixture with MP-TsOH resin that removes DBU and stops acyl migration. This "quenched" distribution allows for a practical isolation of the product with a simple purification.

[0034] Thus, the present invention provides an efficient molecular evolution model in combining the base-catalyzed intramolecular acyl migration of inositol dibenzoate and a boronic acid selector. In the nonlimiting example described

above, it was possible to amplify and accumulate up to 82% of the equilibrium mixture one component, 3,4-di-O-benzoyl-*chiro*-inositol, out of a total of 9 isomers originating from 3,6-di-O-benzoyl-*chiro*-inositol. As a result, 3,6-di-O-benzoyl-*chiro*-inositol, as shown in Figure 2a, efficiently evolved into 3,4-di-O-benzoyl-*chiro*-inositol (Figure 2c) with a boronic acid selector *via* nine mutants (Figure 2b). This is the first application of a boronic acid used to select and amplify a carbohydrate member of a DCL through intramolecular acyl migration.

[0035] The following nonlimiting examples are provided to illustrate the present invention and not to limit it.

[0036] Since it is well known that boronic acids form five-membered cyclic esters preferably with 1,2-*cis* diols in sugar systems,^{12,13} it was envisioned that only one isomer out of the nine, 3,4-dibenzoyl-*chiro*-inositol, which carries two *cis*-vicinal diols, would have a higher binding affinity for two boronic acids, thus stopping further migration. Furthermore, it was also reported that the binding affinity of boronic acid with a sugar depends on the pH of the media.¹³

Materials Used

[0037] All reactions were performed with oven-dried glassware under positive nitrogen pressure. Unless otherwise noted, starting materials and solvents were purchased from Aldrich and Acros organics, and used without further purification. Analytical TLC was carried out on Merck 60 F254 silica gel plates (0.25 mm layer thickness) with visualization by UV light and/or spaying with a 5% solution of phosphomolybdic acid followed by charring with a heat gun. Column chromatography was performed on Merck 60 silica gel (230-400 mesh). ¹H-NMR (200 MHz) spectra were obtained on a Varian Gemini 200 spectrometer. Chemical shifts were reported in parts per million (ppm) relative to the internal standard tetramethylsilane (TMS). Coupling constants (*J*) are in Hertz (Hz). All compounds were identified by LC-MS (Agilent Technology) using a C18 column (4.6×150 mm, 70 minute elution using a solution of 18% CH₃CN-H₂O containing 1% acetic acid), with UV detection at λ=250 nm, and an electrospray ionization source.

General experimental procedure for molecular evolution with pre-equilibrium

[0038] To a solution of **5** (1 mg, 2.6 μmol) in acetonitrile (1 mL) was added DBU (30 μL, 0.20 mmol) at room temperature. After shaking for 1 hour the full migration of 9 isomers was confirmed by HPLC-MS. Phenyl boronic acid was added to the

reaction mixture (10 mg, 0.082 mmol) and was stirred for 7 hours. The distribution change was analyzed by HPLC-MS every hour. For freezing/quenching the distribution, the reaction mixture was filtered through MP-TsOH resin several times to remove DBU.

General experimental procedure for molecular evolution without pre-equilibrium

[0039] To a solution of **5** (1 mg, 2.6 μ mol) in acetonitrile (1 mL) were added phenyl boronic acid (10 mg, 0.082 mmol), and DBU (30 μ L, 0.20 mmol) at room temperature. After stirring for 6 hours, the reaction mixture was analyzed by HPLC-MS.

Synthesis of 1,2,4,5-di-O-isopropylidene-3,6-di-O-benzoyl-chiro-inositol (12**)**

[0040] To a solution of 1,2,4,5-di-O-isopropylidene-3-O-benzoyl-chiro-inositol (**10**)¹¹ (172 mg, 0.47 mmol) in dichloromethane (10 mL) were added pyridine (0.23 mL, 2.8 mmol) and trifluoromethanesulfonic anhydride (0.20 mL, 1.2 mmol) at -15 °C, and the temperature allowed to rise to room temperature. After 2 hours, the reaction mixture was diluted with ethyl acetate, washed with cold 1N HCl solution, saturated NaHCO₃, and brine, dried over MgSO₄, concentrated, and chromatographed on silica gel (EA / Hex = 1 : 10) to give a triflate **11**, which was dissolved in DMF and to which lithium benzoate (120 mg, 0.94 mmol) was added. After stirring at room temperature for 1 day, the reaction mixture was diluted with

ethyl acetate, washed with brine, dried over MgSO_4 , concentrated, and chromatographed on silica gel (EA / Hex = 1 : 10) to give **12** (190 mg, 86.0% in two steps).

Synthesis of 3,6-di-O-benzoyl-chiro-inositol (5)

[0041] A solution of **2** (180 mg, 0.38 mmol) in 80% aqueous AcOH (6 mL) was heated at 100 °C for 2 hours, concentrated under vacuum, and chromatographed on silica gel (EA / Hex = 3 : 1) to give **5** (110 mg, 73%) ^1H NMR (CDCl_3) 3.94-4.08 (2H, m), 4.19 (2H, ddd, $J = 3.6, 7.6, 9.8$), 5.44 (1H, t, $J = 9.8$), 5.53 (1H, t, $J = 3.6$), 7.40-7.58 (6H, m), 7.99-8.10 (4H, m); LC-MS: m/z : 389 $[\text{M}+1]^+$, 371 $[\text{M}-\text{H}_2\text{O}+1]^+$

Synthesis of 3,4-di-O-benzoyl-1,2;5,6-di-O-isopropylidene-chiro-inositol (14)

[0042] To a solution of 1,2;5,6-di-O-isopropylidene-chiro-inositol (**13**)¹¹ (500 mg, 1.9 mmol) in DMF (2 mL) and pyridine (3 mL) was added benzoyl chloride (750 mg, 4.9 mmol). The reaction mixture was stirred for 3 hours at room temperature, diluted with ethyl acetate, washed with water, dried over MgSO_4 , concentrated, and chromatographed on silica gel to give **14** (703 mg, 78.2%).

Synthesis of 3,4-di-O-benzoyl-1,2;5,6-chiro-inositol (7)

[0043] To a solution of **14** (300 mg, 0.64 mmol) in THF (10 mL) and MeOH (2 mL) was added concentrated HCl (0.3 mL). The reaction mixture was stirred for 3 hours. The solvent was removed, and the reaction mixture was diluted with ethyl

acetate, washed with water, dried over MgSO_4 , and concentrated under vacuum. The compound was purified by recrystallization with ethyl acetate and hexane to give **7** (170 mg, 68.4%). $^1\text{H-NMR}$ ($\text{CDCl}_3+\text{MeOH}$) δ 4.15 (2H, t, $J = 2.4$), 4.20 (2H, dd, $J = 4.0$, 6.6), 5.65 (2H, dd, $J = 2.8$, 6.76), 7.31-7.50 (6H, m), 7.92-7.97 (4H, m); LC-MS: m/z : 389 $[\text{M}+1]^+$

[0044] Reaction Conditions: dibenzoyl chiro inositol (**5**) (1 mg), DBU (30 μL , 79 eq.) in acetonitrile (1 mL) with various amounts of phenyl boronic acid

[0045] Reaction conditions: dibenzoyl chiro-inositol (**5**) (1mg), DBU (30 μL , 79 eq.) in acetonitrile (1mL) with phenyl boronic acid (10mg, 32 eq.)

[0046] The foregoing description of the specific embodiments of the present invention will so fully reveal the general nature of the invention that others can, by applying current knowledge, readily modify and/or adapt for various application such specific embodiments without undue experimentation and without departing from the generic concept. Therefore, such adaptations and modifications should and are intended to be comprehended within the meaning and range of equivalents of the disclosed embodiments.

[0047] It is to be understood that the phraseology or terminology employed herein is for the purpose of description

and not of limitation. The means and materials for carrying out disclosed functions may take a variety of alternative forms without departing from the invention. Thus, the expressions "means to..." and "means for..." as may be found in the specification above, and/or in the claims below, followed by a functional statement, are intended to define and cover whatever structural, physical, chemical, or electrical element or structures which may now or in the future exist for carrying out the recited function, whether or not precisely equivalent to the embodiment or embodiments disclosed in the specification above, and it is intended that such expressions be given their broadest interpretation.

References

- (1) (a) Otto, S.; Furlan, R. L. E.; Sanders, J. K. M. *Curr. Opin. Chem. Biol.*, **2002**, *6*, 321-327; (b) Furlan, R. L. E.; Otto, S.; Sanders, J. K. M. *Proc. Natl. Acad. Sci. USA.*, **2002**, *99*, 4801-4804; (c) Rowan, S. J.; Cantrill, S. J.; Cousins, G. R. L.; Sanders, J. K. M.; Stoddart, J. F. *Angew. Chem. Int. Ed.*, **2002**, *41*, 898-952; (d) Ramström, O.; Bunyapaiboonsri, T.; Lohmann, S.; Lehn, J. M. *Biochim. Biophys. Acta*, **2002**, *1572*, 178-186; (e) Lehn, J. M.; Eliseev, A. V. *Science*, **2001**, *291*, 2331-2332; (f) Karan, C.; Miller, B. L. *Drug Discov. Today*, **2000**, *5*, 67-75; (g) Ganesan, A. *Angew. Chem. Int. Ed.*, **1998**, *37*, 2828-2831.
- (2) (a) Brady, P. A.; Bonar-Law, R. P.; Rowan, S. J.; Suckling, C. J.; Sanders, J. K. M. *Chem. Commun.*, **1996**, 319-320; (b) Rowan, S. J.; Brady, P. A.; Sanders, J. K. M. *Angew. Chem. Int. Ed.*, **1996**, *35*, 2143-2145; (c) Rowan, S. J.; Brady, P. A.; Sanders, J. K. M. *Tetrahedron Lett.*, **1996**, *37*, 6013-6016; (d) Rowan, S. J.; Sanders, J. K. M. *J. Org. Chem.*, **1998**, *63*, 1536-1546; (e) Brady, P. A.; Sanders, J. K. M. *J. Chem. Soc., Perkin Trans. 1*, **1997**, 3237-3253; (f) Kaiser, G.; Sanders, J. K. M. *Chem. Commun.*, **2000**, 1763-1764.
- (3) (a) Swann, P. G.; Casanova, R. A.; Desai, A.; Frauenhoff, M. M.; Urbancic, M.; Slomczynska, U.; Hopfinger, A.; Le Breton, G. C.; Venton, D. L. *Biopolymers*, **1996**, *40*, 617-625; (b) Lins, R. J.; Flitsch, S. L.; Turner, N. J.; Irving, E.; Brown, S. A. *Angew. Chem., Int. Ed.*, **2002**, *41*, 3405-3407.
- (4) (a) Roberts, S. L.; Furlan, R. L. E.; Otto, S.; Sanders, J. K. M. *Org. Biomol. Chem.*, **2003**, *1*, 1625-1633; (b) Furlan, R. L. E.; Ng, Y. F.; Cousins, G. R. L.; Redman, J. E.; Sanders, J. K. M. *Tetrahedron*, **2002**, *58*, 771-778; (c) Furlan, R. L. E.; Ng, Y. F.; Otto, S.; Sanders, J. K. M. *J. Am. Chem. Soc.*, **2001**, *123*, 8876-8877; (d) Cousins, G. R. L.; Furlan, R. L. E.; Ng, Y. F.; Redman, J. E.; Sanders, J. K. M. *Angew. Chem., Int. Ed.*, **2001**, *40*, 423-428; (e) Huc, I.; Lehn, J. M. *Proc. Natl. Acad. Sci. USA*, **1997**, *94*, 2106-2110; (f) Star, A.; Goldberg, I.; Fuchs, B. *Angew. Chem., Int. Ed.*, **2000**, *39*, 2685-2689; (g) Storm, O.; Lüning, U. *Chem. Eur. J.* **2002**, *8*, 793-798; (h) Nazarpak-Kandlousy, N.; Zweigenbaum, J.; Henion,

- J.; Eliseev, A. V. *J. Comb. Chem.*, **1999**, *1*, 199-206; (i) Gousins, G. R. L.; Poulsen, S. A.; Sanders, J. K. M. *Chem. Commun.*, **1999**, 1575-1576; (j) Bunyapaiboonsri, T.; Ramström, O.; Lohmann, S.; Lehn, J. M.; Peng, L.; Goeldner, M. *ChemBioChem.*, **2001**, *2*, 438-444; (k) Hochgurtel, M.; Kroth, H.; Piecha, D.; Hofmann, M. W.; Nicolau, C.; Krause, S.; Schaaf, O.; Sonnenmoser, G.; Eliseev, A. *Proc. Natl. Acad. Sci. USA*, **2002**, *99*, 3382-3387; (l) Kuhnert, N.; Rossignolo, G. M.; Lopez-Periago, A. *Org. Biomol. Chem.*, **2003**, *1*, 1157-1170.
- (5) Giger, T.; Wigger, M.; Audetat, S.; Benner, S. A. *Synlett.*, **1998**, 688-692.
- (6) (a) Otto, S.; Furlan, R. L. E.; Sanders, J. K. M. *Science*, **2002**, *297*, 590-593; (b) Otto, S.; Furlan, R. L. E.; Sanders, J. K. M. *J. Am. Chem. Soc.*, **2000**, *122*, 12063-12064; (c) Ramström, O.; Lehn, J. M. *ChemBioChem.*, **2000**, *1*, 41-48; (d) Hioki, H.; Still, W. C. *J. Org. Chem.*, **1998**, *63*, 904-905.
- (7) (a) Eliseev, A. V.; Nelen, M. I. *J. Am. Chem. Soc.*, **1997**, *119*, 1147-1148; (b) Eliseev, A. V.; Nelen, M. I. *Chem. Eur. J.*, **1998**, *4*, 825-834.
- (8) (a) Crego-Calama, M.; Hulst, R.; Fokkens, R.; Nibbering, N. M. M.; Timmerman, P.; Reinhoudt, D. N. *Chem. Commun.*, **1998**, 1021-1022; (b) Crego-Calama, M.; Timmerman, P.; Reinhoudt, D. N. *Angew. Chem., Int. Ed.*, **2000**, *39*, 755-758; (c) Hof, F.; Nuckolls, C.; Rebek, J. J. *Am. Chem. Soc.*, **2000**, *122*, 4251-4252.
- (9) (a) Epstein, D. M.; Choudhary, S.; Churchill, R. M.; Keil, K. M.; Eliseev, A. V.; Morrow, J. R. *Inorg. Chem.*, **2001**, *40*, 1591-1596; (b) Choudhary, S.; Morrow, J. R. *Angew. Chem., Int. Ed.*, **2002**, *41*, 4096-4098; (c) Huc, I.; Krische, M. J.; Funeriu, D. P.; Lehn, J. M. *Eur. J. Inorg. Chem.*, **1999**, 1415-1420; (d) Stulz, E.; Ng, Y. F.; Scott, S. M.; Sanders, J. K. M. *Chem. Commun.*, **2002**, 524-525; (e) Ziegler, M.; Miranda, J. J.; Andersen, U. N.; Johnson, D. W.; Leary, J. A.; Raymond, K. N. *Angew. Chem., Int. Ed.*, **2001**, *40*, 733-736; (f) Albrecht, M.; Blau, O.; Fröhlich, R. *Chem. Eur. J.*, **1999**, *5*, 48-56; (g) Hiraoke, S.; Fujita, M. *J. Am. Chem. Soc.*, **1999**, *121*, 10239-10240; (h) Kubota, Y.; Sakamoto, S.; Yamaguchi, K.; Fujita, M. *Proc. Natl. Acad. Sci. USA*, **2002**, *99*, 4854-4856; (i) Sakai, S.; Shigemasa, Y.; Sasaki, T. *Tetrahedron Lett.*, **1997**, *38*, 8145-8148; (j) Klekota, B.; Hammond, M. H.; Miller, B. L. *Tetrahedron Lett.*, **1997**, *38*, 8639-8642; (k) Klekota, B.; Miller, B. L.

- Tetrahedron*, **1999**, *55*, 11687-11697; (l) Karan, C.; Miller, B. L. *J. Am. Chem. Soc.*, **2001**, *123*, 7455-7456; (m) Case, M. A.; McLendon, G. L. *J. Am. Chem. Soc.*, **2000**, *122*, 8089-8090; (n) Constable, E. C.; Housecroft, C. E.; Kulke, T.; Lazzarini, C.; Schofield, E. R.; Zimmermann, Y. *J. Chem. Soc., Dalton Trans.*, **2001**, 2864-2871; (o) Goral, V.; Nelen, M. I.; Eliseev, A. V.; Lehn, J. M. *Proc. Natl. Acad. Sci. USA*, **2001**, *98*, 1347-1352.
- (10) (a) Chung, S. K.; Chang, Y. T. *J. Chem. Soc., Chem. Commun.*, **1995**, 13-14; (b) Chung, S. K.; Chang, Y. T.; Ryu, Y. *Pure App. Chem.*, **1996**, *68*, 931-935.
- (11) For synthesis of inositol derivatives, refer to supplementary information, and (a) Khersonsky, S. M.; Chang, Y. T. *Carbohydr. Res.*, **2002**, *337*, 75-78; (b) Falshaw, A.; Hart, J. B.; Tyler, P. C. *Carbohydr. Res.*, **2000**, *329*, 301-308.
- (12) (a) James, T. D.; Sandanayake, S.; Shinkai, S. *Angew. Chem., Int. Ed.*, **1996**, *35*, 1910-1922; (b) Wiecko, J.; Sherman, W. R. *J. Am. Chem. Soc.*, **1979**, *101*, 979-983.
- (13) (a) Eggert, H.; Frederiksen, J.; Morin, C.; Norrild, J. C. *J. Org. Chem.*, **1999**, *64*, 3846-3852; (b) Arimori, S.; Bell, M. L.; Oh, C. S.; Frimat, K. A.; James, T. D. *Chem. Commun.*, **2001**, 1836-1837; (c) Arimori, S.; Ushiroda, S.; Peter, L. M.; Jenkins, A. T. A.; James, T. D. *Chem. Commun.*, **2002**, 2368-2369; (d) Yang, W.; He, H.; Drueckhammer, D. G. *Angew. Chem., Int. Ed.*, **2001**, *40*, 1714-1718; (e) Cabell, L. A.; Monahan, M. K.; Anslyn, E. V. *Tetrahedron Lett.*, **1999**, *40*, 7753-7756; (f) James, T. D.; Sandanayake, S.; Iguchi, R.; Shinkai, S. *J. Am. Chem. Soc.*, **1995**, *117*, 8982-8987.

WHAT IS CLAIMED IS:

1. A method for generating a dynamic carbohydrate library comprising subjecting a carbohydrate scaffold to intramolecular acyl migration using a boronic acid selector.
2. The method according to claim 1 wherein the carbohydrate scaffold is a monomeric carbohydrate.
3. The method according to claim 1 wherein the carbohydrate scaffold is selected from the group consisting of dimmers and oligomers having monomers connected by an ester bond.
4. The method according to claim 1 wherein the boronic acid selector is phenyl boronic acid.
5. The method according to claim 1 wherein the generating is conducted in a basic medium.
6. The method according to claim 1 wherein the carbohydrate scaffold is inositol.
7. The method according to claim 7 wherine the inositol is *chiro*-inositol.

ABSTRACT OF THE DISCLOSURE:

An efficient molecular evolution system was obtained by combining two novel approaches in the field, intramolecular acyl migration for DCL generation and boronic acid as a selector. 1,4-Di-O-benzoyl chiro-inositol in acetonitrile with DBU as a base generated a reversible equilibrium of 9 regioisomers. Only 3,4-di-O-benzoyl chiro-inositol, containing two sets of cis-diols, is most favored toward binding with phenyl boronic acid, and its percentage of the DCL composition was amplified by up to 82%.

FIGURE 1




FIGURE 2




FIGURE 3






FIGURE 4



Effect of phenyl boronic acid on the amplification of 7

FIGURE 5



Time dependent amplification of 7

FIGURE 6





FIGURE 7 a


LC data (250 nm)



LC diagram of compound 7

FIGURE 7 b


Mass data



MS diagram of compound 7

FIGURE 8 a


LC data (250 nm)



LC diagram of compound 5


FIGURE 8 b

Mass data




MS diagram of compound 5

FIGURE 9



$$5 \times 5 \times 5 \times 5 \times 6 = 3750 \text{ isomers}$$

FIGURE 10



TOC Intramolecular Acyl Migration of *chiro*-Inositol dibenzoate: Overwhelming Selection by Phenyl Boronic Acid